arXiv:1209.5234v1 [math.CA] 24 Sep 2012

UMD BANACH SPACES AND THE MAXIMAL REGULARITY FOR THE HERMITE, BESSEL AND LAGUERRE OPERATORS

VÍCTOR ALMEIDA, JORGE J. BETANCOR, AND ALEJANDRO J. CASTRO

Abstract. In this paper we prove that the maximal regularity property for Cauchy problems associated with the Hermite, Bessel and Laguerre operators on $L^p(\Omega,\mathbb{B})$, $1<p<\infty$ characterizes the UMD property for the Banach space $\mathbb{B}$. Here, $\Omega=\mathbb{R}$ in the Hermite setting and $\Omega=(0,\infty)$ in the Bessel and Laguerre contexts.

1. Introduction

Suppose that $\mathbb{B}$ is a complex Banach space, A is a (possible unbounded) operator on $\mathbb{B}$ such that $D(A)$ is dense in $\mathbb{B}$. We consider the Cauchy problem

$$\left\{\begin{array}{l} u'(t)+Au(t)=f(t), \quad 0\leq t<T,\\ u(0)=0, \end{array}\right. \tag{1}$$

where $f:(0,T)\to\mathbb{B}$ is measurable and $0<T\leq\infty$.

Let $1<p<\infty$ and $0<T\leq\infty$. We say that the operator A has the maximal L^p-regularity property on the interval $(0,T)$ when for every $f\in L^p((0,T),\mathbb{B})$ there exists a unique solution $u\in L^p((0,T),\mathbb{B})$ of (1), and

$$\|u'\|_{L^p((0,T),\mathbb{B})}+\|Au\|_{L^p((0,T),\mathbb{B})}\leq C\|f\|_{L^p((0,T),\mathbb{B})},$$

for a certain $C>0$ which does not depend on f. This property of maximal regularity is important, for instance, in nonlinear problems and a lot of authors have studied it in the last years ([3], [4], [5], [6], [7], [8] and [29]).

Assume that $-A$ generates a C_0-semigroup $\{T_t\}_{t>0}$ on $\mathbb{B}$. Then the unique mild solution of (1) is defined by

$$u(t)=\int_0^t T_{t-s}(f(s))ds, \quad t\in(0,T),$$

for every $f\in L^p((0,T),\mathbb{B})$.

If A is bounded, it has maximal L^p-regularity on $(0,T)$ for every $1<p<\infty$ and $0<T<\infty$, and if also there exists $M,m>0$ such that $\|T_t\|\leq Me^{-mt}$, $t>0$, A has maximal L^p-regularity on $(0,\infty)$ for each $1<p<\infty$. When $\{T_t\}_{t>0}$ is analytic and A has maximal L^p-regularity on $(0,T)$ for some $T>0$ and $1<p<\infty$, A has maximal L^q-regularity on $(0,T)$, for every $1<q<\infty$. Moreover, if $1<p<\infty$ and there exists maximal L^p-regularity for A on $(0,T_0)$ for some $T_0\in(0,\infty)$, then A has maximal L^p-regularity on $(0,T)$ for every $T>0$.

Le Merdy [27, Theorem 2.4] gave an example of an operator A generating a bounded analytic semigroup such that A has maximal L^p-regularity on $(0,T)$ for every $T>0$, and it has not maximal L^p-regularity on $(0,\infty)$, with $1<p<\infty$.

2010 *Mathematics Subject Classification.* 47D06, 35B65, 42B25.

Key words and phrases. UMD spaces, maximal regularity, Hermite, Bessel and Laguerre operators.

The authors are partially supported by MTM2010/17974. The third author is also supported by a FPU grant from the Government of Spain.

It can be actually a not easy problem to determinate whether an operator A satisfies the maximal L^p-regularity property. The very beginning work of Grisvard [23] is remarkable about this question.

The Banach spaces with the UMD property play an important role in the study of maximal L^p-regularity for operators. It is well known that the Hilbert transform $\mathcal{H}$ defined, for every $f \in L^p(\mathbb{R})$, $1 < p < \infty$,

$$\mathcal{H}(f)(x) = \lim_{\epsilon \to 0^+} \frac{1}{\pi} \int_{|x-y|>\epsilon} \frac{f(y)}{x-y} dy, \quad a.e.\ x \in \mathbb{R},$$

is bounded from $L^p(\mathbb{R})$ into itself, $1 < p < \infty$. If X is a Banach space, $\mathcal{H}$ is extended to $L^p \otimes X$, $1 < p < \infty$, in a natural way. We say that X is UMD when the Hilbert transform can be extended from $L^p(\mathbb{R}) \otimes X$ to $L^p(\mathbb{R}, X)$ as a bounded operator from $L^p(\mathbb{R}, X)$ into itself for some (equivalently, for any) $1 < p < \infty$. The main properties of UMD Banach spaces can be encountered in [16], [17] and [36].

Dore and Venni [20, Theorem 3.2] established the maximal L^p-regularity for invertible operators A on UMD-Banach spaces $\mathbb{B}$ such that, for every $\gamma \in \mathbb{R} \setminus \{0\}$, the imaginary power $A^{i\gamma}$ of A is bounded and $\|A^{i\gamma}\| \leq Ke^{\theta|\gamma|}$, where $K > 0$ and $0 < \theta < \pi/2$. It is currently an open problem to know whether this condition on the imaginary power is necessary to have maximal L^p-regularity.

Brezis (see [18]) posed the following question: given a Banach space $\mathbb{B}$, is it true that every negative generator $-A$ of a bounded analytic semigroup on $\mathbb{B}$ has the maximal L^p-regularity? When $\mathbb{B}$ is a Hilbert space the answer of this question is affirmative (see [19, Lemma 3.1, p. 211]), but it is unknown under which conditions on the Banach space $\mathbb{B}$ this property holds.

In the sequel we say that a semigroup of operators generated by $-A$ has the maximal L^p-regularity provided that A has it.

As it is well-known if S is a positive bounded operator from $L^p(\Omega, \mu)$ into itself, where $1 \leq p < \infty$ and (Ω, μ) is a measure space, the operator $S \otimes I_X$ can be extended, for every Banach space X, to the Lebesgue-Bochner space $L^p(\Omega, \mu, X)$ as a bounded operator, that we continue denoting by S, from $L^p(\Omega, \mu, X)$ into itself.

For every $t > 0$, we represent, as usual, by $P_t(f)$ the classical Poison integral of $f \in L^p(\mathbb{R})$, $1 \leq p < \infty$, that is,

$$P_t(f)(x) = \int_{\mathbb{R}} P_t(x-y) f(y) dy, \quad x \in \mathbb{R},$$

where

$$P_t(x) = \frac{1}{\pi} \frac{t}{t^2 + x^2}, \quad x \in \mathbb{R}.$$

Thus, P_t is a positive operator and $P_t \otimes I_X$ can be extended to $L^2(\mathbb{R}, X)$, for every $t > 0$.

A partial answer for Brezis' question was given by Coulhon and Lamberton [18, p. 160] who established the following result.

Theorem A. *Let X be a Banach space. Then, the Poisson semigroup $\{P_t\}_{t>0}$ on $\mathbb{B} = L^2(\mathbb{R}, X)$ has maximal L^p-regularity on $(0, \infty)$, if, and only if, X is UMD.*

Note that $L^2(\mathbb{R}, X)$ is UMD provided that X is UMD ([36, Proposition 3, (*ii*)]). Moreover, according to [24, Theorem, p. 402], if X is not UMD then $\left(-\frac{d^2}{dx^2}\right)^{i\gamma}$, $\gamma \in \mathbb{R} \setminus \{0\}$, is not bounded in $L^2(\mathbb{R}, X)$.

Our objective in this paper is to establish results analogous to Theorem A when the classical Poisson semigroup is replaced by the ones associated with Bessel, Hermite and Laguerre operators.

We consider for every $\alpha > 0$ the Bessel operator

$$\Delta_\alpha = -x^{-\alpha}\frac{d}{dx}x^{2\alpha}\frac{d}{dx}x^{-\alpha} = -\frac{d^2}{dx^2} + \frac{\alpha(\alpha-1)}{x^2}, \quad x \in (0,\infty).$$

If J_ν denotes the Bessel function of first kind and order ν, we have that

$$\Delta_\alpha\left[\sqrt{xy}J_{\alpha-1/2}(xy)\right] = y^2\sqrt{xy}J_{\alpha-1/2}(xy), \quad x \in (0,\infty).$$

According to [33, (16.4)] the Poisson kernel for Δ_α is the following

$$P_t^{\Delta_\alpha}(x,y) = \frac{2\alpha(xy)^\alpha t}{\pi}\int_0^\pi \frac{(\sin\theta)^{2\alpha-1}}{[|x-y|^2+t^2+2xy(1-\cos\theta)]^{\alpha+1}}d\theta, \quad t,x,y \in (0,\infty),$$

and the Poisson integral $P_t^{\Delta_\alpha}(f)$ of $f \in L^p(0,\infty)$, $1 \le p \le \infty$, is defined by

$$P_t^{\Delta_\alpha}(f)(x) = \int_0^\infty P_t^{\Delta_\alpha}(x,y)f(y)dy, \quad x \in (0,\infty),\ t > 0.$$

The Bessel-Poisson semigroup $\{P_t^{\Delta_\alpha}\}_{t>0}$ is positive and contractive in $L^p(0,\infty)$, $1 \le p \le \infty$. Harmonic analysis associated to Bessel operators was initiated by Muckenhoupt and Stein ([33]). In the last years, this Bessel harmonic analysis has been developed in a Banach valued setting (see [10] and [13]).

The Hermite operator on $\mathbb{R}$ is defined by $H = -\frac{1}{2}(\frac{d^2}{dx^2} - x^2)$. We have that, for every $k \in \mathbb{N}$,

$$Hh_k = \left(k + \frac{1}{2}\right)h_k,$$

where h_k denotes the k-th Hermite function given by

$$h_k(x) = (\sqrt{k}2^k k!)^{-1/2}H_k(x)e^{-x^2/2}, \quad x \in \mathbb{R},$$

and H_k represents the k-th Hermite polynomial [39, pp. 105–106]. The operator $-H$ generates the positive semigroup of contractions $\{W_t^H\}_{t>0}$ in $L^p(\mathbb{R})$, being, for every $f \in L^p(\mathbb{R})$, $1 \le p \le \infty$,

$$W_t^H(f)(x) = \int_{\mathbb{R}} W_t^H(x,y)f(y)dy, \quad x \in \mathbb{R} \text{ and } t > 0,$$

and for each $x, y \in \mathbb{R}$, $t > 0$,

$$W_t^H(x,y) = \left(\frac{e^{-t}}{\pi(1-e^{-2t})}\right)^{1/2}\exp\left[-\frac{1}{2}\frac{1+e^{-2t}}{1-e^{-2t}}(x^2+y^2) + \frac{2e^{-t}}{1-e^{-2t}}xy\right].$$

The Poisson semigroup $\{P_t^H\}_{t>0}$ associated with the Hermite operator is defined, for every $f \in L^p(\mathbb{R})$, $1 \le p \le \infty$, as

$$P_t^H(f)(x) = \int_{\mathbb{R}} P_t^H(x,y)f(y)dy, \quad x \in \mathbb{R} \text{ and } t > 0,$$

where, by using the subordination formula we can write

$$P_t^H(x,y) = \frac{t}{\sqrt{4\pi}}\int_0^\infty \frac{e^{-t^2/4u}}{u^{3/2}}W_u^H(x,y)du, \quad x,y \in \mathbb{R} \text{ and } t > 0.$$

Muckenhoupt studied in [30] and [31] the harmonic analysis for the Hermite operator in dimension one. In the last years, Thangavelu [40], Stempak and Torrea [37] and [38] have investigated it in higher dimensions. Banach valued harmonic analysis operators in the Hermite context have been studied in [2] and [14].

We consider the Laguerre differential operator L_α, $\alpha > 0$, on $(0,\infty)$ defined by

$$L_\alpha = \frac{1}{2}\left(-\frac{d^2}{dx^2} + x^2 + \frac{\alpha(\alpha-1)}{x^2}\right), \quad x \in (0,\infty).$$

We have that

$$L_\alpha \varphi_k^\alpha = \left(2k + \alpha + \frac{1}{2}\right)\varphi_k^\alpha,$$

where

$$\varphi_k^\alpha(x) = \left(\frac{2\Gamma(k+1)}{\Gamma(k+\alpha+1/2)}\right)^{1/2} e^{-x^2/2} x^\alpha \ell_k^{\alpha-1/2}(x^2), \quad x \in (0,\infty),$$

and, for every $k \in \mathbb{N}$, ℓ_k^α represents the k-th Laguerre polynomial of order α ([39, pp. 100–102]).

The operator $-L_\alpha$ generates the positive semigroup of contractions $\{W_t^{L_\alpha}\}_{t>0}$ in $L^p(0,\infty)$, where, for every $f \in L^p(0,\infty)$, $1 \le p \le \infty$,

$$W_t^{L_\alpha}(f)(x) = \int_0^\infty W_t^{L_\alpha}(x,y) f(y)\,dy, \quad x \in (0,\infty),\ t > 0,$$

being, for every $t, x, y \in (0,\infty)$,

$$W_t^{L_\alpha}(x,y) = \frac{2\sqrt{xy}e^{-t}}{1-e^{-2t}} I_{\alpha-1/2}\left(\frac{2xye^{-t}}{1-e^{-2t}}\right) \exp\left[-\frac{1}{2}\frac{1+e^{-2t}}{1-e^{-2t}}(x^2+y^2)\right].$$

By using the subordination formula we can write the Poisson kernel $P_t^{L_\alpha}(x,y)$, $t, x, y \in (0,\infty)$, associated to L_α as follows

$$P_t^{L_\alpha}(x,y) = \frac{t}{\sqrt{4\pi}} \int_0^\infty \frac{e^{-t^2/4u}}{u^{3/2}} W_u^{L_\alpha}(x,y)\,du, \quad t, x, y \in (0,\infty).$$

The Poisson semigroup for the Laguerre operator L_α is defined by

$$P_t^{L_\alpha}(f)(x) = \int_0^\infty P_t^{L_\alpha}(x,y) f(y)\,dy, \quad t, x \in (0,\infty),$$

for every $f \in L^p(0,\infty)$, $1 \le p \le \infty$.

The starting point for the harmonic analysis in Laguerre context are the works of Muckenhoupt [31] and [32]. In the last years we remark the results of Nowak and Stempak [34] and [35] in higher dimensions. Laguerre harmonic analysis in a Banach valued setting has been developed in [14] and [15].

The main result of this paper is the following.

Theorem 1.1. *Let X be a Banach space and $\alpha > 0$. The following assertions are equivalent.*

- (a) *X is UMD.*
- (b) *The Bessel Poisson semigroup $\{P_t^{\Delta_\alpha}\}_{t>0}$ on $L^2((0,\infty), X)$ has maximal L^p-regularity on $(0,\infty)$.*
- (c) *The Hermite Poisson semigroup $\{P_t^H\}_{t>0}$ on $L^2(\mathbb{R}, X)$ has maximal L^p-regularity on $(0,\infty)$.*
- (d) *The Laguerre Poisson semigroup $\{P_t^{L_\alpha}\}_{t>0}$ on $L^2((0,\infty), X)$ has maximal L^p-regularity on $(0,\infty)$.*

According to [9, Theorem 1.2] the imaginary powers $H^{i\gamma}$ (respectively, $\Delta_\alpha^{i\gamma}$ and $L_\alpha^{i\gamma}$), are not bounded in $L^p(\mathbb{R}, X)$ (respectively, in $L^p((0,\infty), X)$), $1 < p < \infty$, when the Banach space is not UMD.

In order to prove the maximal L^p-regularity for the bounded analytic semigroup of operators $\{T_t\}_{t>0}$ generated by the operator $-A$ on a Banach space $\mathbb{B}$ we consider the operator $K(\{T_u\}_{u>0})$ defined by

$$K(\{T_u\}_{u>0})f(t) = \int_0^t \partial_t T_{t-s}(f(s))ds, \quad t > 0. \tag{2}$$

Since $\|\partial_t T_t\|_{L(\mathbb{B})} \leq C/t$, $t > 0$, where $L(\mathbb{B})$ denotes the space of bounded operators from $\mathbb{B}$ into itself, the convergence of the integral in (2) depends on the properties of f. Suppose that $f \in C_c^1((0,\infty), D(A))$, that is, $f : (0,\infty) \longrightarrow D(A)$ is a C^1-function with compact support. Then, we have that

$$\begin{aligned}\int_0^t \|\partial_t T_{t-s}(f(s))\|_{\mathbb{B}}\, ds \leq & C\left(\int_0^t \frac{\|f(s)-f(t)\|_{\mathbb{B}}}{t-s}ds + t\|Af(t)\|_{\mathbb{B}}\right)\\ \leq & Ct\left(\|f'\|_{L^\infty((0,\infty),\mathbb{B})} + \|Af(t)\|_{\mathbb{B}}\right), \quad t > 0.\end{aligned}$$

The subspace $C_c^1(0,\infty) \otimes D(A)$ is dense in $L^p((0,\infty),\mathbb{B})$, $1 < p < \infty$, because $D(A)$ is dense in $\mathbb{B}$. The semigroup $\{T_t\}_{t>0}$ of operators (or the operator A) has the maximal L^p-regularity property if, and only if, the operator $K(\{T_u\}_{u>0})$ can be extended from $C_c^1(0,\infty) \otimes D(A)$ to $L^q((0,\infty);\mathbb{B})$ as a bounded operator from $L^q((0,\infty);\mathbb{B})$ into itself, for some (equivalently, for every) $1 < q < \infty$. For the sake of simplicity, when the operator $K(\{T_u\}_{u>0})$ satisfies this property we say that it is bounded from $L^q((0,\infty);\mathbb{B})$ into itself.

If $\{T_t\}_{t>0}$ denotes Bessel, Hermite or Laguerre Poisson semigroups, to prove Theorem 1.1 we will show that the operator $K(\{T_u\}_{u>0})$ is bounded from $L^2((0,\infty), L^2(\Omega, X))$ into itself if, and only if, the operator $K(\{P_u\}_{u>0})$ is bounded from $L^2((0,\infty), L^2(\mathbb{R}, X))$ into itself. Here $\Omega = (0,\infty)$ when Bessel or Laguerre operators are considered and $\Omega = \mathbb{R}$ when we work with the Hermite operator. Actually, we split the operator $K(\{T_u\}_{u>0})$ in two parts that we call local and global parts

$$K(\{T_u\}_{u>0}) = K^{loc}(\{T_u\}_{u>0}) + K^{glob}(\{T_u\}_{u>0}).$$

The operator $K^{glob}(\{T_u\}_{u>0})$ is bounded in $L^2((0,\infty), L^2(\Omega, X))$ for every Banach space X. On the other hand, the operator $K^{loc}(\{T_u\}_{u>0})$ is bounded in $L^2((0,\infty), L^2(\Omega, X))$ if, and only if, $K(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty), L^2(\mathbb{R}, X))$. Then, according to Theorem 1.1, the $L^2((0,\infty), L^2(\Omega, X))$-boundedness of $K(\{T_u\}_{u>0})$ characterizes the UMD property for X.

In Sections 2, 3 and 4 we prove Theorem 1.1 in the Bessel, Hermite and Laguerre setting, respectively.

Throughout this paper by $C, c > 0$ we always denote positive constants that can change in each occurrence.

2. Proof of Theorem 1.1 for the Bessel operator

In this section we prove de equivalence $(a) \Leftrightarrow (b)$ in Theorem 1.1.

We denote by $C_c^\infty(\Omega)$ the space of smooth functions in Ω with compact support, where $\Omega = \mathbb{R} \setminus \{0\}$ or $\Omega = (0,\infty)$. It is well-known that $C_c^\infty(0,\infty)$ is a dense subspace of $L^2(0,\infty)$ and $C_c^\infty(\mathbb{R} \setminus \{0\})$ is dense in $L^2(\mathbb{R})$.

We consider the operators $K(\{P_u^{\Delta_\alpha}\}_{u>0})$ and $K(\{P_u\}_{u>0})$ defined, for each $t, x \in (0,\infty)$, by

$$K(\{P_u^{\Delta_\alpha}\}_{u>0})(f)(t,x) = \int_0^t \partial_t P_{t-s}^{\Delta_\alpha}(f(s,\cdot))(x)ds, \quad f \in C_c^\infty(0,\infty) \otimes C_c^\infty(0,\infty) \otimes X,$$

and for $t \in (0,\infty)$, $x \in \mathbb{R}$,

$$K(\{P_u\}_{u>0})(f)(t,x) = \int_0^t \partial_t P_{t-s}(f(s,\cdot))(x)ds, \quad f \in C_c^\infty(0,\infty) \otimes C_c^\infty(\mathbb{R} \setminus \{0\}) \otimes X.$$

Note that, for every $f \in C_c^\infty(0,\infty) \otimes C_c^\infty(0,\infty) \otimes X$,

$$\left(\int_0^t \partial_t P_{t-s}(f(s,\cdot))ds\right)(x) = \int_0^t \partial_t P_{t-s}(f(s,\cdot))(x)ds, \quad \text{a.e. } x \in \mathbb{R},$$

where the first integral is understood in the $L^2(\mathbb{R},X)$-Bochner sense and the integral in the right hand side is understood in the X-Bochner sense. Analogous properties hold when Hermite, Bessel and Laguerre Poisson semigroups replace the classical Poisson semigroup.

We will prove that $K(\{P_u^{\Delta_\alpha}\}_{u>0})$ is bounded from $L^2((0,\infty)^2,X)$ into itself if, and only if, $K(\{P_u\}_{u>0})$ is bounded from $L^2((0,\infty)\times\mathbb{R},X)$ into itself. Then, by using Theorem A we deduce $(a) \Leftrightarrow (b)$ in Theorem 1.1.

We define the operator $K_+(\{P_u\}_{u>0})$, for each $t,x \in (0,\infty)$, as follows

$$K_+(\{P_u\}_{u>0})(f)(t,x) = \int_0^t \int_0^\infty \partial_t P_{t-s}(x-y)f(s,y)dyds, \quad f \in C_c^\infty(0,\infty) \otimes C_c^\infty(0,\infty) \otimes X.$$

In the first step we reduce the boundedness of $K(\{P_u\}_{u>0})$ in $L^2((0,\infty)\times\mathbb{R},X)$ to the boundedness of $K_+(\{P_u\}_{u>0})$ in $L^2((0,\infty)^2,X)$.

Lemma 2.1. *Let X be a Banach space. Then, $K(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$ if, and only if, $K_+(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)^2,X)$.*

Proof. Let $f \in C_c^\infty(0,\infty) \otimes C_c^\infty(0,\infty) \otimes X$. We defined

$$f_0(t,x) = \begin{cases} f(t,x), & t,x>0,\\ 0, & t>0,\ x \le 0. \end{cases} \tag{3}$$

It is clear that $f_0 \in C_c^\infty(0,\infty) \otimes C_c^\infty(\mathbb{R}\setminus\{0\}) \otimes X$ and

$$K_+(\{P_u\}_{u>0})(f)(t,x) = K(\{P_u\}_{u>0})(f_0)(t,x), \quad t,x \in (0,\infty).$$

Then, if $K(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$, $K_+(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)^2,X)$.

Suppose now that $K_+(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)^2,X)$ and let $f \in C_c^\infty(0,\infty)\otimes C_c^\infty(\mathbb{R}\setminus\{0\}) \otimes X$. We can write for every $t>0$ and $x \in \mathbb{R}$,

$$\begin{aligned} K(\{P_u\}_{u>0})(f)(t,x) &= \int_0^t \int_{\mathbb{R}} \partial_t P_{t-s}(x-y)f(s,y)dyds\\ &= \int_0^t \int_0^\infty \partial_t P_{t-s}(x-y)f(s,y)dyds + \int_0^t \int_0^\infty \partial_t P_{t-s}(x+y)f(s,-y)dyds, \end{aligned}$$

and

$$\begin{aligned} \int_0^\infty \int_{\mathbb{R}} \|K(\{P_u\}_{u>0})(f)(t,x)\|_X^2 dxdt &= \int_0^\infty \int_0^\infty \|K(\{P_u\}_{u>0})(f)(t,x)\|_X^2 dxdt\\ &\quad + \int_0^\infty \int_0^\infty \|K(\{P_u\}_{u>0})(f)(t,-x)\|_X^2 dxdt\\ &\le 2\Big[|K_+(\{P_u\}_{u>0})(f_+)\|^2_{L^2((0,\infty)^2,X)} + \|K_+(\{P_u\}_{u>0})(f_-)\|^2_{L^2((0,\infty)^2,X)}\\ &\quad + \|K_-(\{P_u\}_{u>0})(f_+)\|^2_{L^2((0,\infty)^2,X)} + \|K_-(\{P_u\}_{u>0})(f_-)\|^2_{L^2((0,\infty)^2,X)}\Big], \end{aligned} \tag{4}$$

where $f_+(t,x) = f(t,x)$, $f_-(t,x) = f(t,-x)$, $t,x \in (0,\infty)$, and,

$$K_-(\{P_u\}_{u>0})(g)(t,x) = \int_0^t \int_0^\infty \partial_t P_{t-s}(x+y)g(s,y)dyds, \quad g \in C_c^\infty(0,\infty) \otimes C_c^\infty(0,\infty) \otimes X.$$

The operator K_- is bounded from $L^2((0,\infty)^2,X)$ into itself. Indeed, we have that

$$|\partial_t P_{t-s}(x+y)| = \frac{1}{\pi}\left|\frac{|t-s|^2-(x+y)^2}{(|t-s|^2+(x+y)^2)^2}\right| \leq \frac{1}{\pi}\frac{1}{|t-s|^2+(x+y)^2}, \quad s,t,x,y\in(0,\infty).$$

Hence,

$$\begin{aligned}\|K_-(\{P_u\}_{u>0})(g)(t,x)\|_X &\leq \frac{1}{\pi}\int_0^t\int_0^\infty \frac{1}{|t-s|^2+(x+y)^2}\|g(s,y)\|_X dyds\\ &\leq \frac{1}{\pi}\int_0^\infty\left(\int_0^x \frac{1}{|t-s|^2+x^2}\|g(s,y)\|_X dy + \int_x^\infty \frac{1}{|t-s|^2+y^2}\|g(s,y)\|_X dy\right)ds\\ &\leq \frac{1}{\pi}\left(\frac{1}{x}\int_0^x P_*(\|\tilde{g_0}(\cdot,y)\|_X)(t)dy + \int_x^\infty \frac{1}{y}P_*(\|\tilde{g_0}(\cdot,y)\|_X)(t)dy\right), \quad t,x\in(0,\infty),\end{aligned} \tag{5}$$

where

$$\tilde{g_0}(t,y) = \begin{cases} g_0(t,y), & t,y\in(0,\infty),\\ 0 & t\in(-\infty,0),\ y\in(0,\infty),\end{cases} \tag{6}$$

and P_* represents the maximal operator defined by

$$P_*(h) = \sup_{u>0}|P_u(h)|, \quad h\in L^2(\mathbb{R}).$$

According to [26, p. 244, (9.9.1) and (9.9.2)], the Hardy operators H_0 and H_∞ given by

$$H_0(h)(x) = \frac{1}{x}\int_0^x h(y)dy, \quad x\in(0,\infty),$$

and

$$H_\infty(h)(x) = \int_x^\infty \frac{h(y)}{y}dy, \quad x\in(0,\infty),$$

are bounded in $L^2(0,\infty)$. Also, the maximal operator P_* is bounded in $L^2(\mathbb{R})$. Then, from (5) we deduce that K_- is bounded in $L^2((0,\infty)^2,X)$.

By (4) it follows that $K(\{P_u\}_{u>0})$ is bounded from $L^2((0,\infty)\times\mathbb{R},X)$ into itself provided that $K_+(\{P_u\}_{u>0})$ is bounded from $L^2((0,\infty)^2,X)$ into itself. □

We now consider, for every $t,x\in(0,\infty)$ and $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$, the following "local" and "global" operators

- $K_+^{loc}(\{P_u\}_{u>0})f(t,x) = \int_0^t\int_{x/2}^{2x}\partial_t P_{t-s}(x-y)f(s,y)dyds,$
- $K_+^{glob}(\{P_u\}_{u>0})f(t,x) = K_+(\{P_u\}_{u>0})f(t,x) - K_+^{loc}(\{P_u\}_{u>0})f(t,x),$
- $K^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0})f(t,x) = \int_0^t\int_{x/2}^{2x}\partial_t P_{t-s}^{\Delta_\alpha}(x,y)f(s,y)dyds,$

and

- $K^{glob}(\{P_u^{\Delta_\alpha}\}_{u>0})f(t,x) = K(\{P_u^{\Delta_\alpha}\}_{u>0})f(t,x) - K^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0})f(t,x).$

We are going to see that the "global" operators are bounded in $L^2((0,\infty)^2,X)$.

Lemma 2.2. *Let X be a Banach space. Then, the operators $K_+^{glob}(\{P_u\}_{u>0})$ and $K^{glob}(\{P_u^{\Delta_\alpha}\}_{u>0})$ are bounded from $L^2((0,\infty)^2,X)$ into itself.*

Proof. Let $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$. The argument used in the proof of Lemma 2.1 allows us to get, for every $t,x\in(0,\infty)$,

$$\|K_+^{glob}(\{P_u\}_{u>0})(f)(t,x)\|_X \leq C\left(\frac{1}{x}\int_0^{x/2} P_*(\|\tilde{f}_0(\cdot,y)\|_X)(t)dy + \int_{2x}^\infty \frac{1}{y}P_*(\|\tilde{f}_0(\cdot,y)\|_X)(t)dy\right),$$

where the extension $\tilde{f}_0$ of f is defined as in (6). Then, K_+^{glob} is bounded in $L^2((0,\infty)^2, X)$.

On the other hand, we have that

$$\begin{aligned}\partial_t P_{t-s}^{\Delta_\alpha}(x,y) =& \frac{2\alpha(xy)^\alpha}{\pi}\left\{\int_0^\pi \frac{(\sin\theta)^{2\alpha-1}}{[|x-y|^2+|t-s|^2+2xy(1-\cos\theta)]^{\alpha+1}}d\theta\right.\\ &\left.-2(\alpha+1)|t-s|^2\int_0^\pi \frac{(\sin\theta)^{2\alpha-1}}{[|x-y|^2+|t-s|^2+2xy(1-\cos\theta)]^{\alpha+2}}d\theta\right\},\ s,t,x,y\in(0,\infty).\end{aligned}$$

According to [33, p. 86, (b)] we obtain

$$\begin{aligned}\left|\partial_t P_{t-s}^{\Delta_\alpha}(x,y)\right| \leq& C(xy)^\alpha\int_0^\pi \frac{(\sin\theta)^{2\alpha-1}}{[|x-y|^2+|t-s|^2+2xy(1-\cos\theta)]^{\alpha+1}}d\theta\\ \leq& C\frac{1}{|x-y|^2+|t-s|^2},\quad s,t,x,y\in(0,\infty).\end{aligned}$$

Then, as above we can prove that $K^{glob}(\{P_u^{\Delta_\alpha}\}_{u>0})$ is bounded in $L^2((0,\infty)^2, X)$. □

Hence, we will finish the proof once we show that $K_+^{loc}(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)^2, X)$ if, and only if, $K^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0})$ is bounded in $L^2((0,\infty)^2, X)$. We collect this property in the next Lemma.

Lemma 2.3. *Let X be a Banach space. Then, the operator D^{loc} given by*

$$D^{loc} = K^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0}) - K_+^{loc}(\{P_u\}_{u>0}),$$

is bounded in $L^2((0,\infty)^2, X)$.

Proof. Let $f \in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$. We split the operator $K^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0})$ as follows

$$K^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0}) = K_1^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0}) + K_2^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0}),$$

being,

$$K_1^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0})(f)(t,x) = \int_0^t\int_{x/2}^{2x}\partial_t P_{t-s}^{\Delta_\alpha,1}(x,y)f(s,y)dyds,\quad t,x\in(0,\infty),$$

and

$$P_t^{\Delta_\alpha,1}(x,y) = \frac{2\alpha(xy)^\alpha}{\pi}\int_0^{\pi/2}\frac{t(\sin\theta)^{2\alpha-1}}{[|x-y|^2+t^2+2xy(1-\cos\theta)]^{\alpha+1}}d\theta,\quad t,x,y\in(0,\infty).$$

By defining

$$P_t^{\Delta_\alpha,2}(x,y) = P_t^{\Delta_\alpha}(x,y) - P_t^{\Delta_\alpha,1}(x,y),\quad t,x,y\in(0,\infty),$$

we can write

$$\begin{aligned}\left|\partial_t P_{t-s}^{\Delta_\alpha,2}(x,y)\right| \leq& C(xy)^\alpha\int_{\pi/2}^\pi \frac{(\sin\theta)^{2\alpha-1}}{[|x-y|^2+|t-s|^2+2xy(1-\cos\theta)]^{\alpha+1}}d\theta\\ \leq& C\frac{1}{|t-s|^2+x^2},\quad s,t,x\in(0,\infty),\ x/2<y<2x.\end{aligned}$$

Hence, we get

$$\|K_2^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0})(f)(t,x)\|_X \leq \frac{C}{x}\int_{x/2}^{2x}P_*(\|\tilde{f}_0(\cdot,y)\|_X)(t)dy,\quad t,x\in(0,\infty).$$

Then, since the operator $\mathcal{L}$ defined by

$$\mathcal{L}(h)(x) = \frac{1}{x}\int_{x/2}^{2x}h(y)dy,\quad x\in(0,\infty),$$

is bounded in $L^2(0,\infty)$, we conclude that $K_2^{loc}(\{P_u^{\Delta_\alpha}\}_{u>0})$ is bounded in $L^2((0,\infty)^2, X)$.

We consider the kernels

$$P_t^{\Delta_\alpha,1,1}(x,y)=\frac{2\alpha(xy)^\alpha}{\pi}\int_0^{\pi/2}\frac{t\theta^{2\alpha-1}}{[|x-y|^2+t^2+2xy(1-\cos\theta)]^{\alpha+1}}d\theta,\quad t,x,y\in(0,\infty),$$

$$P_t^{\Delta_\alpha,1,2}(x,y)=\frac{2\alpha(xy)^\alpha}{\pi}\int_0^{\pi/2}\frac{t\theta^{2\alpha-1}}{[|x-y|^2+t^2+xy\theta^2]^{\alpha+1}}d\theta,\quad t,x,y\in(0,\infty).$$

and we define the operators

$$D_1^{loc}(f)(t,x)=\int_0^t\int_{x/2}^{2x}\partial_t\left[P_{t-s}^{\Delta_\alpha,1}(x,y)-P_{t-s}^{\Delta_\alpha,1,1}(x,y)\right]f(s,y)dyds,\quad t,x\in(0,\infty)$$

and

$$D_2^{loc}(f)(t,x)=\int_0^t\int_{x/2}^{2x}\partial_t\left[P_{t-s}^{\Delta_\alpha,1,1}(x,y)-P_{t-s}^{\Delta_\alpha,1,2}(x,y)\right]f(s,y)dyds,\quad t,x\in(0,\infty).$$

In order to estimate the kernels of D_1^{loc} and D_2^{loc} we proceed as in [11, pp. 482–485]. By using mean value theorem, we get for every $s,t,x\in(0,\infty)$ and $x/2<y<2x$,

$$\begin{aligned}\Big|\partial_t&\left[P_{t-s}^{\Delta_\alpha,1}(x,y)-P_{t-s}^{\Delta_\alpha,1,1}(x,y)\right]\Big|+\Big|\partial_t\left[P_{t-s}^{\Delta_\alpha,1,1}(x,y)-P_{t-s}^{\Delta_\alpha,1,2}(x,y)\right]\Big|\\&\leq C(xy)^\alpha\int_0^{\pi/2}\frac{\theta^{2\alpha+1}}{[|x-y|^2+|t-s|^2+xy\theta^2]^{\alpha+1}}d\theta\\&\leq C\frac{(xy)^\alpha}{[|x-y|^2+|t-s|^2+xy]^{\alpha+1}}\left[1+\log_+\left(\frac{xy}{|x-y|^2+|t-s|^2}\right)\right].\end{aligned}$$

Hence, for $j=1,2$,

$$\begin{aligned}\|D_j^{loc}(f)(t,x)\|_X\leq&C\int_0^t\int_{x/2}^{2x}\frac{1}{x^2+|t-s|^2}\left(1+\log_+\frac{x}{|x-y|}\right)\|f(s,y)\|_Xdyds\\\leq&C\frac{1}{x}\int_{x/2}^{2x}\left(1+\log_+\frac{x}{|x-y|}\right)P_*(\|\tilde{f}_0(\cdot,y)\|_X)(t)dy,\quad t,x\in(0,\infty).\end{aligned}$$

Since

$$\frac{1}{x}\int_{x/2}^{2x}\left(1+\log_+\frac{x}{|x-y|}\right)dy=\int_{1/2}^2\left(1+\log_+\frac{1}{|1-z|}\right)dz,\quad x\in(0,\infty),$$

Jensen's inequality allows us to show that the operator $\mathcal{A}$ defined by

$$\mathcal{A}(g)(x)=\frac{1}{x}\int_{x/2}^{2x}\left(1+\log_+\frac{x}{|x-y|}\right)g(y)dy,\quad x\in(0,\infty),$$

is bounded from $L^2(0,\infty)$ into itself. Hence, the operators D_1^{loc} and D_2^{loc} are bounded in $L^2((0,\infty)^2,X)$.

Finally, we consider the operator D_3^{loc} as follows

$$D_3^{loc}(f)(t,x)=\int_0^t\int_{x/2}^{2x}\partial_t\left[P_{t-s}^{\Delta_\alpha,1,2}(x,y)-P_{t-s}(x-y)\right]f(s,y)dyds,\quad t,x\in(0,\infty).$$

As in [11, p. 486] we can write, fore every $t,x,y\in(0,\infty)$,

$$P_t^{\Delta_\alpha,1,2}(x,y)=P_t(x-y)-\frac{2\alpha(xy)^\alpha}{\pi}\int_{\pi/2}^\infty\frac{t\theta^{2\alpha-1}}{[|x-y|^2+t^2+xy\theta^2]^{\alpha+1}}d\theta.$$

Then, we get

$$D_3^{loc}(f)(t,x)=\int_0^t\int_{x/2}^{2x}\partial_tP_{t-s}^{\Delta_\alpha,1,3}(x,y)f(s,y)dyds,\quad t,x\in(0,\infty),$$

where

$$P_t^{\Delta_\alpha,1,3}(x,y)=-\frac{2\alpha(xy)^\alpha}{\pi}\int_{\pi/2}^\infty\frac{t\theta^{2\alpha-1}}{[|x-y|^2+t^2+xy\theta^2]^{\alpha+1}}d\theta,\quad t,x,y\in(0,\infty).$$

We have that

$$\begin{aligned}\left|\partial_t P^{\Delta_\alpha,1,3}_{t-s}(x,y)\right| &\leq C\int_{\pi/2}^{\infty}\frac{(xy)^\alpha\theta^{2\alpha-1}}{[|x-y|^2+|t-s|^2+xy\theta^2]^{\alpha+1}}d\theta\\ &\leq C(xy)^\alpha\left[\frac{1}{[|x-y|^2+|t-s|^2+xy\pi^2/4]^{\alpha+1}}+\int_{\pi^2/4}^{\infty}\frac{xyu^\alpha}{[|x-y|^2+|t-s|^2+xyu]^{\alpha+2}}du\right]\\ &\leq C(xy)^\alpha\left[\frac{1}{[|t-s|^2+xy]^{\alpha+1}}+(xy)^{-\alpha+1}\int_{\pi^2/4}^{\infty}\frac{1}{[|t-s|^2+xyu]^2}du\right]\\ &\leq C\frac{1}{|t-s|^2+x^2},\quad s,t,x\in(0,\infty)\text{ and } x/2<y<2x.\end{aligned}$$

We can write

$$\|D_3^{loc}(f)(t,x)\|_X\leq C\int_0^t\int_{x/2}^{2x}\frac{\|f(s,y)\|_X}{|t-s|^2+x^2}dyds\leq C\frac{1}{x}\int_{x/2}^{2x}P_*(\|\tilde{f}_0(\cdot,y)\|_X)(t)dy,\ t,x\in(0,\infty).$$

Hence, D_3^{loc} is bounded from $L^2((0,\infty)^2,X)$ into itself. We conclude that D^{loc} is bounded in $L^2((0,\infty)^2,X)$. □

3. PROOF OF THEOREM 1.1 FOR THE HERMITE OPERATOR

In this section we prove the equivalence $(a)\Leftrightarrow(c)$ in Theorem 1.1.

We consider the operator defined by

$$K\left(\{P_u^H\}_{u>0}\right)(f)(t,x)=\int_0^t\partial_t P_{t-s}^H(f(s,\cdot))(x)ds,\quad x\in\mathbb{R}\text{ and } t>0,$$

for every $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(\mathbb{R})\otimes X$.

We denote by ρ the following function

$$\rho(x)=\begin{cases}1/2, & |x|\leq 1,\\ \dfrac{1}{1+|x|}, & |x|>1.\end{cases}$$

This function ρ plays an important role in the study of harmonic analysis associated to the Hermite operator (see [12] and [22]).

We split the operator $K\left(\{P_u^H\}_{u>0}\right)$ as follows

$$K\left(\{P_u^H\}_{u>0}\right)=K^{loc}\left(\{P_u^H\}_{u>0}\right)+K^{glob}\left(\{P_u^H\}_{u>0}\right),$$

where

$$K^{loc}\left(\{P_u^H\}_{u>0}\right)(f)(t,x)=\int_0^t\int_{|x-y|<\rho(x)}\partial_t P_{t-s}^H(x,y)f(s,y)dyds,\quad x\in\mathbb{R}\text{ and } t>0,$$

for every $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(\mathbb{R})\otimes X$.

In a similar way we decompose the operator $K\left(\{P_u\}_{u>0}\right)$ on $C_c^\infty(0,\infty)\otimes C_c^\infty(\mathbb{R})\otimes X$ by

$$K\left(\{P_u\}_{u>0}\right)=K^{loc}\left(\{P_u\}_{u>0}\right)+K^{glob}\left(\{P_u\}_{u>0}\right).$$

In a first step, we are going to see that the operator $K^{glob}\left(\{P_u^H\}_{u>0}\right)$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$.

Lemma 3.1. *Let X be a Banach space. Then, the operator $K^{glob}\left(\{P_u^H\}_{u>0}\right)$ is bounded from $L^2((0,\infty)\times\mathbb{R},X)$ into itself.*

Proof. Let $f \in C_c^\infty(0,\infty) \otimes C_c^\infty(\mathbb{R}) \otimes X$. We have that

$$\begin{aligned}\left|\partial_t P^H_{t-s}(x,y)\right| \leq& C\int_0^\infty \frac{e^{-|t-s|^2/4u}}{u^{3/2}}\left(1+\frac{|t-s|^2}{u}\right)W^H_u(x,y)du\\ \leq& C\int_0^\infty \frac{e^{-c|t-s|^2/u}}{u^{3/2}}W^H_u(x,y)du, \quad x,y\in\mathbb{R} \text{ and } s,t\in(0,\infty).\end{aligned}$$

Then,

$$\begin{aligned}&\left\|K^{glob}\left(\{P^H_u\}_{u>0}\right)(f)(t,x)\right\|_X\\ &\quad\leq C\int_0^t\int_{|x-y|\geq\rho(x)}\left(\int_0^{\rho(x)^2}+\int_{\rho(x)^2}^\infty\right)\frac{e^{-c|t-s|^2/u}}{u^{3/2}}W^H_u(x,y)du\|f(s,y)\|_X dyds\\ &\quad= C\left[K^{glob}_{H,1}(\|f\|_X)(t,x)+K^{glob}_{H,2}(\|f\|_X)(t,x)\right], \quad x\in\mathbb{R} \text{ and } t\in(0,\infty). \qquad (7)\end{aligned}$$

Let $h \in L^2((0,\infty)\times\mathbb{R})$. We detone by h_0 the following extension of h

$$h_0(t,x)=\begin{cases} h(t,x), & t>0,\ x\in\mathbb{R},\\ 0, & t\leq 0,\ x\in\mathbb{R}.\end{cases}$$

Since

$$W^H_t(x,y)\leq Ce^{-ct}W_{ct}(x-y), \quad x,y\in\mathbb{R} \text{ and } t>0, \tag{8}$$

where $W_t(z)=e^{-|z|^2/4t}/\sqrt{4\pi t}$, $z\in\mathbb{R}$ and $t>0$, represents the classical heat semigroup, we can write

$$\begin{aligned}\left|K^{glob}_{H,1}(h)(t,x)\right| \leq& C\int_0^t\int_{|x-y|\geq\rho(x)}\int_0^{\rho(x)^2}\frac{e^{-c|t-s|^2/u}}{u^{3/2}}\frac{e^{-c|x-y|^2/u}}{u^{1/2}}|h(s,y)|dudyds\\ \leq& C\int_{|x-y|\geq\rho(x)}\int_0^{\rho(x)^2}\frac{e^{-c|x-y|^2/u}}{u^{3/2}}W_*(|h_0(\cdot,y)|)(t)dudy, \quad x\in\mathbb{R} \text{ and } t>0,\end{aligned}$$

where $W_*(g)=\sup_{u>0}|W_u(g)|$. We get

$$\begin{aligned}\left|K^{glob}_{H,1}(h)(t,x)\right| \leq& C\int_0^{\rho(x)^2}\int_{|x-y|\geq\rho(x)}\frac{e^{-c\rho(x)^2/u}}{u}\frac{e^{-c|x-y|^2/u}}{u^{1/2}}W_*(|h_0(\cdot,y)|)(t)dydu\\ \leq C&\left(\int_0^{\rho(x)^2}\frac{du}{\rho(x)^2}\right)W_*\left[W_*(|h_0|)(t)\right](x)\leq CW_*\left[W_*(|h_0|)(t)\right](x), \quad x\in\mathbb{R} \text{ and } t>0.\end{aligned}$$

Since the maximal operator W_* is bounded in $L^2(\mathbb{R})$ we deduce that the operator $K^{glob}_{H,1}$ is bounded from $L^2((0,\infty)\times\mathbb{R})$ into itself.

In order to show that $K^{glob}_{H,2}$ is bounded in $L^2((0,\infty)\times\mathbb{R})$ we will use the following estimate

$$W^H_t(x,y)\leq C\frac{e^{-c|x-y|^2/t}}{\sqrt{t}}\left(\frac{\rho(x)^2}{t}\right)^2, \quad x,y\in\mathbb{R} \text{ and } t>0. \tag{9}$$

Indeed, we have that

$$|x-y|^2+|x+y|^2\geq x^2\geq\frac{C}{\rho(x)^2}, \quad |x|>1.$$

Then, since $\rho(x)=1/2$, $|x|\leq 1$,

$$W^H_t(x,y)\leq Ce^{-c(t+|x+y|^2+|x-y|^2)}\leq C\frac{e^{-c|x-y|^2/t}}{\sqrt{t}}\left(\frac{\rho(x)^2}{t}\right)^2, \quad x,y\in\mathbb{R} \text{ and } t>1.$$

Suppose now that $0<t\leq 1$. If $|x|\leq 1$ and $y\in\mathbb{R}$,

$$W^H_t(x,y)\leq C\frac{e^{-c(t+|x-y|^2/t)}}{\sqrt{t}}\leq C\frac{e^{-c|x-y|^2/t}}{\sqrt{t}}\left(\frac{\rho(x)^2}{t}\right)^2.$$

Moreover, if $|x|>1$ and $y\in\mathbb{R}$,

$$\begin{aligned}W_t^H(x,y) \le& C\frac{e^{-c(t+|x-y|^2/t+t|x+y|^2)}}{\sqrt{t}} \le C\frac{e^{-c|x-y|^2/t}e^{-ctx^2}}{\sqrt{t}} \le C\frac{e^{-c|x-y|^2/t}e^{-ct/\rho(x)^2}}{\sqrt{t}}\\ \le& C\frac{e^{-c|x-y|^2/t}}{\sqrt{t}}\left(\frac{\rho(x)^2}{t}\right)^2, \quad xy\ge 0,\end{aligned}$$

and

$$\begin{aligned}W_t^H(x,y) \le& C\frac{e^{-c(t+|x-y|^2/t)}}{\sqrt{t}} \le C\frac{e^{-ct}e^{-c|x-y|^2/t}e^{-cx^2/t}}{\sqrt{t}} \le C\frac{e^{-c|x-y|^2/t}}{\sqrt{t}}\left(\frac{1}{tx^2}\right)^2\\ \le& C\frac{e^{-c|x-y|^2/t}}{\sqrt{t}}\left(\frac{\rho(x)^2}{t}\right)^2, \quad xy\le 0.\end{aligned}$$

Thus, (9) is established.

We can write

$$\begin{aligned}\left|K_{H,2}^{glob}(h)(t,x)\right| \le& C\int_0^t\int_{|x-y|\ge\rho(x)}\int_{\rho(x)^2}^{\infty}\frac{e^{-c|t-s|^2/u}}{u^{1/2}}\frac{e^{-c|x-y|^2/u}}{u^{3/2}}\frac{\rho(x)^4}{u^2}|h(s,y)|dudyds\\ \le& C\int_{|x-y|\ge\rho(x)}\left(\int_{\rho(x)^2}^{\infty}\frac{\rho(x)^4}{u^{7/2}}e^{-c|x-y|^2/u}du\right)W_*(|h_0(\cdot,y)|)(t)dy\\ \le& C\int_{|x-y|\ge\rho(x)}\left(\rho(x)^6\int_1^{\infty}\frac{e^{-c|x-y|^2/v\rho(x)^2}}{(v\rho(x)^2)^{7/2}}dv\right)W_*(|h_0(\cdot,y)|)(t)dy\\ \le& C\int_{|x-y|\ge\rho(x)}\left(\frac{1}{\rho(x)}\int_1^{\infty}\left(\frac{\rho(x)}{|x-y|}\right)^3\frac{dv}{v^2}\right)W_*(|h_0(\cdot,y)|)(t)dy\\ \le& C\frac{1}{\rho(x)}\int_{|x-y|\ge\rho(x)}\left(\frac{\rho(x)}{|x-y|}\right)^3W_*(|h_0(\cdot,y)|)(t)dy\\ =& C\frac{1}{\rho(x)}\sum_{k=0}^{\infty}\int_{2^k\rho(x)\le|x-y|<2^{k+1}\rho(x)}\left(\frac{\rho(x)}{|x-y|}\right)^3W_*(|h_0(\cdot,y)|)(t)dy\\ \le& C\sum_{k=0}^{\infty}\frac{1}{2^{3k}\rho(x)}\int_{|x-y|<2^{k+1}\rho(x)}W_*(|h_0(\cdot,y)|)(t)dy\\ \le& C\mathcal{M}\left[W_*(|h_0|)(t)\right](x), \quad x\in\mathbb{R} \text{ and } t>0.\end{aligned}$$

Here $\mathcal{M}$ represents the Hardy-Littlewood maximal function. Since W_* and $\mathcal{M}$ are bounded operators in $L^2(\mathbb{R})$, it follows that the operator $K_{H,2}^{glob}$ is bounded from $L^2((0,\infty)\times\mathbb{R})$ into itself.

From (7) we conclude that $K^{glob}(\{P_u^H\}_{u>0})$ is bounded from $L^2((0,\infty)\times\mathbb{R},X)$ into itself. □

In the second step we show that the operator $\mathcal{D}^{loc}$ defined by

$$\mathcal{D}^{loc}=K^{loc}(\{P_u^H\}_{u>0})-K^{loc}(\{P_u\}_{u>0})$$

is bounded in $L^2((0,\infty)\times\mathbb{R},X)$.

Lemma 3.2. *Let X be a Banach space. Then, the operator $\mathcal{D}^{loc}$ is bounded from $L^2((0,\infty)\times\mathbb{R},\ X)$ into itself.*

Proof. Let $f \in C_c^\infty(0,\infty) \otimes C_c^\infty(\mathbb{R}) \otimes X$. The subordination formula and (8) allow us to write

$$\begin{aligned}\left\|\mathcal{D}^{loc}(f)(t,x)\right\|_X \leq & C\Big(\int_0^t \int_{|x-y|<\rho(x)} \int_0^{\rho(x)^2} \frac{e^{-c|t-s|^2/u}}{u^{3/2}} \left|W_u^H(x,y) - W_u(x-y)\right| \|f(s,y)\|_X dudyds \\ & + \int_0^t \int_{|x-y|<\rho(x)} \int_{\rho(x)^2}^\infty \frac{e^{-c|t-s|^2/u}}{u^{3/2}} W_{cu}(x-y)\|f(s,y)\|_X dudyds\Big) \\ = & C\left[\mathcal{D}_1^{loc}(\|f\|_X)(t,x) + \mathcal{D}_2^{loc}(\|f\|_X)(t,x)\right], \quad x \in \mathbb{R} \text{ and } t \in (0,\infty).\end{aligned}$$

We now prove that $\mathcal{D}_j^{loc}$, $j=1,2$, is a bounded operator from $L^2((0,\infty)\times\mathbb{R})$ into itself. Let $h \in L^2((0,\infty)\times\mathbb{R})$. We have that

$$\begin{aligned}\left|\mathcal{D}_2^{loc}(h)(t,x)\right| \leq & C \int_{|x-y|<\rho(x)} \left(\int_{\rho(x)^2}^\infty \frac{du}{u^{3/2}}\right) W_*(|h_0(\cdot,y)|)(t)dy \\ \leq & \frac{C}{\rho(x)} \int_{|x-y|<\rho(x)} W_*(|h_0(\cdot,y)|)(t)dy \\ \leq & C\mathcal{M}\left[W_*(|h_0|)(t)\right](x), \quad x \in \mathbb{R} \text{ and } t>0.\end{aligned}$$

Hence, $\mathcal{D}_2^{loc}$ is bounded in $L^2((0,\infty)\times\mathbb{R})$.

In order to analyze $\mathcal{D}_1^{loc}$ we need to show that

$$(10) \qquad \left|W_t^H(x,y) - W_t(x-y)\right| \leq C\frac{t}{\rho(x)^2} W_{ct}(x-y), \quad |x-y|<\rho(x) \text{ and } 0<t<\rho(x)^2.$$

Indeed, by using Kato-Trotter's formula and (8) we get, for every $x,y \in \mathbb{R}$ and t>0,

$$\begin{aligned}&\left|W_t^H(x,y) - W_t(x-y)\right| = \int_0^t \int_{\mathbb{R}} W_s(x-z) W_{t-s}^H(z,y) z^2 dzds \\ &\leq C\int_0^t \int_{\mathbb{R}} W_s(x-z) W_{c(t-s)}(z-y) z^2 dzds \\ &\leq C\left(\int_0^{t/2} \int_{\mathbb{R}} W_s(x-z) W_{c(t-s)}(z-y) z^2 dzds + \int_0^{t/2} \int_{\mathbb{R}} W_{t-s}(x-z) W_{cs}(z-y) z^2 dzds\right).\end{aligned}$$

On the other hand, we have that

$$\begin{aligned}\int_{\mathbb{R}} e^{-c|x-z|^2/s} z^2 dz = & \sqrt{s}\int_{\mathbb{R}} e^{-w^2}(x^2 + sw^2 - 2xw\sqrt{s})dw \\ \leq & C\sqrt{s}\int_{\mathbb{R}} e^{-w^2}(x^2 + w^2 + xw)dw, \quad x \in \mathbb{R},\ 0<s<1.\end{aligned}$$

If $|x| \leq 1$, then

$$\int_{\mathbb{R}} e^{-w^2}(x^2+w^2+xw)dw \leq \int_{\mathbb{R}} e^{-w^2}(1+w^2+w)dw \leq C \leq \frac{C}{\rho(x)^2},$$

and, when $|x| \geq 1$,

$$\begin{aligned}\int_{\mathbb{R}} e^{-w^2}(x^2+w^2+xw)dw \leq & \int_{\mathbb{R}} e^{-w^2}\left(\frac{1}{\rho(x)^2} + w^2 + \frac{w}{\rho(x)}\right)dw \\ \leq & \frac{C}{\rho(x)^2}\int_{\mathbb{R}} e^{-w^2}(1+w^2+w)dw \leq \frac{C}{\rho(x)^2},\end{aligned}$$

because $\rho(x) \leq 1$. Hence,

$$\int_{\mathbb{R}} e^{-c|x-z|^2/s} z^2 dz \leq C\frac{\sqrt{s}}{\rho(x)^2}, \quad x \in \mathbb{R} \text{ and } 0<s<1.$$

It follows that

$$\begin{aligned}
\left|W_t^H(x,y)-W_t(x-y)\right| &\leq C\Big(\int_0^{t/2}\int_{\mathbb{R}}\frac{e^{-c[|x-z|^2/s+|z-y|^2/(t-s)]}}{s^{1/2}(t-s)^{1/2}}z^2dzds\\
&+\int_0^{t/2}\int_{\mathbb{R}}\frac{e^{-c[|x-z|^2/(t-s)+|z-y|^2/s]}}{s^{1/2}(t-s)^{1/2}}z^2dzds\Big)\\
&\leq C\int_0^{t/2}\int_{\mathbb{R}}\frac{e^{-c[|x-z|^2+|z-y|^2]/t}}{t^{1/2}}\left(\frac{e^{-c|x-z|^2/s}}{s^{1/2}}+\frac{e^{-c|y-z|^2/s}}{s^{1/2}}\right)z^2dzds\\
&\leq C\frac{e^{-c|x-y|^2/t}}{t^{1/2}}\int_0^{t/2}\int_{\mathbb{R}}\left(\frac{e^{-c|x-z|^2/s}}{s^{1/2}}+\frac{e^{-c|y-z|^2/s}}{s^{1/2}}\right)z^2dzds\\
&\leq C\frac{t}{\rho(x)^2}W_{ct}(x-y),\quad |x-y|<\rho(x)\text{ and }0<t<\rho(x)^2,
\end{aligned}$$

because $\rho(x)/C\leq\rho(y)\leq C\rho(x)$, provided that $|x-y|<\rho(x)$. Thus (10) is proved.

By using (10) we deduce that

$$\begin{aligned}
\left|\mathcal{D}_1^{loc}(h)(t,x)\right| \leq & C\int_0^t\int_{|x-y|<\rho(x)}|h(s,y)|\int_0^{\rho(x)^2}\frac{e^{-c|t-s|^2/u}}{u^{1/2}\rho(x)^2}\frac{e^{-c|x-y|^2/u}}{u^{1/2}}dudyds\\
\leq & C\int_{|x-y|<\rho(x)}\left(\int_0^{\rho(x)^2}\frac{du}{u^{1/2}\rho(x)^2}\right)W_*(|h_0(\cdot,y)|)(t)dy\\
\leq & \frac{C}{\rho(x)}\int_{|x-y|<\rho(x)}W_*(|h_0(\cdot,y)|)(t)dy\\
\leq & C\mathcal{M}\left[W_*(|h_0|)(t)\right](x),\quad x\in\mathbb{R}\text{ and }t>0.
\end{aligned}$$

Hence, $\mathcal{D}_1^{loc}$ is bounded from $L^2((0,\infty)\times\mathbb{R})$ into itself. We conclude that $\mathcal{D}^{loc}$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$. □

Note that the operators $K^{glob}(\{P_u^H\}_{u>0})$ and $\mathcal{D}^{loc}$ are bounded in $L^2((0,\infty)\times\mathbb{R},X)$ for every Banach space X.

By Lemmas 3.1 and 3.2 the operator $K(\{P_u^H\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$ if, and only if, the operator $K^{loc}(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$. According to Theorem A our equivalence ($(a)\Leftrightarrow(c)$ in Theorem 1.1) will be showed once we establish that $K(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$ if, and only if, $K^{loc}(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$. In order to do this we use some ideas developed in [1] and [25].

Lemma 3.3. *Let X be a Banach space. Then, the operator $K(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$ if, and only if, $K^{loc}(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$.*

Proof. Suppose that $K(\{P_u\}_{u>0})$ is bounded from $L^2((0,\infty)\times\mathbb{R},X)$ into itself. Let $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(\mathbb{R})\otimes X$. There exists $C_0>0$ such that

$$\frac{1}{C_0}\rho(x)\leq\rho(y)\leq C_0\rho(x),\quad |x-y|\leq\rho(x).$$

According to [21, Proposition 5] (see also [22, Lemma 2.3]) we choose a sequence $(x_k)_{k=1}^\infty\subseteq\mathbb{R}$ such that

(a) $\mathbb{R}=\bigcup_{k=1}^\infty B(x_k,\rho(x_k))$.

(b) For every $M>0$ there exists $m\in\mathbb{N}$ such that, for every $\ell\in\mathbb{N}$,

$$\#\{k\in\mathbb{N}:B(x_k,M\rho(x_k))\cap B(x_\ell,M\rho(x_\ell))\neq\varnothing\}\leq m,$$

where $\#A$ denotes the cardinal of the set $A\subseteq\mathbb{N}$.

To simplify notation we write $I_k = B(x_k, \rho(x_k))$ and $MI_k = B(x_k, M\rho(x_k))$, $M > 0$, $k \in \mathbb{N}$.

We define the operator T as follows

$$T(f)(t,x) = \sum_{k=1}^{\infty} \chi_{I_k}(x) \int_0^t \int_{(C_0+1)I_k} \partial_t P_{t-s}(x-y) f(s,y) dy ds, \quad x \in \mathbb{R} \text{ and } t > 0.$$

We have that

$$\begin{aligned} \|T(f)\|^2_{L^2((0,\infty)\times\mathbb{R},X)} \leq & C \sum_{k=1}^{\infty} \left\| K(\{P_u\}_{u>0}) \left(\chi_{(C_0+1)I_k}(y) f(s,y) \right) \right\|^2_{L^2((0,\infty)\times\mathbb{R},X)} \\ \leq & C \sum_{k=1}^{\infty} \left\| \chi_{(C_0+1)I_k}(y) f(s,y) \right\|^2_{L^2((0,\infty)\times\mathbb{R},X)} \\ \leq & C \|f\|^2_{L^2((0,\infty)\times\mathbb{R},X)} . \end{aligned}$$

To see that $K^{loc}(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$ it is enough to show that the difference operator $\mathcal{D} = T - K^{loc}(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$.

If $x \in I_k$ and $|x-y| < \rho(x)$, then $y \in (C_0+1)I_k$. Also, if $x \in I_k$ and $y \in (C_0+1)I_k$, then

$$|x-y| \leq \rho(x_k) + (C_0+1)\rho(x_k) \leq C_0(C_0+2)\rho(x).$$

Hence, we have that

$$\left\{(x,y) \in \mathbb{R}^2 : |x-y| < \rho(x)\right\} \subseteq \bigcup_{k\in\mathbb{N}} \left\{(x,y) \in \mathbb{R}^2 : x \in I_k,\ y \in (C_0+1)I_k\right\},$$

and

$$\begin{aligned} \bigcup_{k\in\mathbb{N}} \left\{(x,y) \in \mathbb{R}^2 : x \in I_k,\ y \in (C_0+1)I_k\right\} \setminus \left\{(x,y) \in \mathbb{R}^2 : |x-y| < \rho(x)\right\} \\ \subseteq \left\{(x,y) \in \mathbb{R}^2 : \rho(x) \leq |x-y| \leq C_0(C_0+2)\rho(x)\right\}. \end{aligned}$$

We can write

$$\begin{aligned} \mathcal{D}(f)(t,x) = & \int_0^t \int_{\mathbb{R}} \left[\sum_{k=1}^{\infty} \chi_{I_k}(x)\chi_{(C_0+1)I_k}(y) - \chi_{\{(x,y)\in\mathbb{R}^2 : |x-y|<\rho(x)\}}(x,y) \right] \\ & \times \partial_t P_{t-s}(x-y) f(s,y) dy ds, \quad x \in \mathbb{R} \text{ and } t > 0. \end{aligned}$$

Hence, we get

$$\begin{aligned} \|\mathcal{D}(f)(t,x)\|_X \leq & C \int_0^t \int_{\rho(x)\leq|x-y|\leq C_0(C_0+2)\rho(x)} |\partial_t P_{t-s}(x-y)| \, \|f(s,y)\|_X dy ds \\ \leq & C \int_0^t \int_{\rho(x)\leq|x-y|\leq C_0(C_0+2)\rho(x)} \frac{1}{|t-s|^2 + |x-y|^2} \|f(s,y)\|_X dy ds \\ \leq & C \int_0^t \int_{\rho(x)\leq|x-y|\leq C_0(C_0+2)\rho(x)} \frac{1}{|t-s|^2 + \rho(x)^2} \|f(s,y)\|_X dy ds \\ \leq & \frac{C}{\rho(x)} \int_{|x-y|\leq C_0(C_0+2)\rho(x)} P_* \left(\|f_0(\cdot,y)\|_X \right)(t) dy \\ \leq & C\mathcal{M}\left[P_* \left(\|f_0\|_X \right)(t) \right](x), \quad x \in \mathbb{R} \text{ and } t > 0. \end{aligned}$$

We conclude that $\mathcal{D}$ is bounded from $L^2((0,\infty)\times\mathbb{R},X)$ into itself. Then, $K^{loc}(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$.

Assume now that $K^{loc}(\{P_u\}_{u>0})$ is a bounded operator from $L^2((0,\infty)\times\mathbb{R},X)$ into itself. We are going to see that $K(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$. Let $f \in C_c^\infty(0,\infty) \otimes C_c^\infty(\mathbb{R}) \otimes X$.

For every $R>0$ we define $f_R(t,x)=f(t,Rx)$ and $f^R(t,x)=f(Rt,x)$, $x\in\mathbb{R}$ and $t>0$. Straightforward manipulations allows us to show that

$$K(\{P_u\}_{u>0})(f)(t,x)=K(\{P_u\}_{u>0})((f_R)^R)\left(\frac{t}{R},\frac{x}{R}\right),\quad x\in\mathbb{R}\text{ and } t,R>0.$$

Suppose that $f(t,x)=0$, $|x|>a$, $t>0$, where $a>0$. Fix $b>0$. We have that, for every $R>0$,

$$\rho\left(\frac{x}{R}\right)=\begin{cases}1/2, & |x|\le R,\\ \dfrac{R}{R+|x|}>\dfrac{R}{R+b}, & |x|>R,\end{cases}\qquad |x|\le b,$$

and

$$\left|y-\frac{x}{R}\right|\le\frac{a+b}{R},\quad |y|\le\frac{a}{R}\text{ and } |x|\le b.$$

Hence, there exists $R_0>0$ such that $|y-x/R|<\rho(x/R)$, $|x|\le b$, $|y|\le a/R$ and $R\ge R_0$. Hence,

$$\begin{aligned}K(\{P_u\}_{u>0})(f)(t,x)=&\frac{1}{\pi}\int_0^{t/R}\int_{-a/R}^{a/R}\frac{(t/R-s)^2-(x/R-y)^2}{\left[(t/R-s)^2+(x/R-y)^2\right]^2}(f_R)^R(s,y)dyds\\ =&\frac{1}{\pi}\int_0^{t/R}\int_{|y-x/R|<\rho(x/R)}\frac{(t/R-s)^2-(x/R-y)^2}{\left[(t/R-s)^2+(x/R-y)^2\right]^2}(f_R)^R(s,y)dyds\\ =&K^{loc}(\{P_u\}_{u>0})((f_R)^R)\left(\frac{t}{R},\frac{x}{R}\right),\quad |x|\le b,\ t>0\text{ and } R\ge R_0.\end{aligned}$$

Since $K^{loc}(\{P_u\}_{u>0})$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$, it follows that

$$\begin{aligned}\int_0^\infty\int_{-b}^b\|K(\{P_u\}_{u>0})(f)(t,x)\|_X^2\,dxdt=&\int_0^\infty\int_{\mathbb{R}}\left\|K^{loc}(\{P_u\}_{u>0})((f_{R_0})^{R_0})\left(\frac{t}{R_0},\frac{x}{R_0}\right)\right\|_X^2dxdt\\ =&R_0^2\int_0^\infty\int_{\mathbb{R}}\left\|K^{loc}(\{P_u\}_{u>0})((f_{R_0})^{R_0})\,(t,x)\right\|_X^2dxdt\\ \le&CR_0^2\int_0^\infty\int_{\mathbb{R}}\|f\,(R_0t,R_0x)\|_X^2\,dxdt\\ \le&C\|f\|^2_{L^2((0,\infty)\times\mathbb{R},X)}.\end{aligned}$$

Hence,

$$\|K(\{P_u\}_{u>0})(f)\|^2_{L^2((0,\infty)\times\mathbb{R},X)}\le C\|f\|^2_{L^2((0,\infty)\times\mathbb{R},X)}.$$

□

4. PROOF OF THEOREM 1.1 FOR LAGUERRE OPERATORS

We are going to prove the equivalence $(a)\Leftrightarrow(d)$ in Theorem 1.1.

We consider the operator

$$K\left(\{P_u^{L_\alpha}\}_{u>0}\right)(f)(t,x)=\int_0^t\partial_tP_{t-s}^{L_\alpha}(f(s,\cdot))(x)ds,\quad t,x\in(0,\infty),$$

for every $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$.

Since we showed in Section 3, $(a)\Leftrightarrow(c)$ in Theorem 1.1, it is enough to prove that $K\left(\{P_u^{L_\alpha}\}_{u>0}\right)$ is bounded in $L^2((0,\infty)^2,X)$ if, and only if, $K\left(\{P_u^{H}\}_{u>0}\right)$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$.

We consider the operator

$$K_+\left(\{P_u^{H}\}_{u>0}\right)(f)(t,x)=\int_0^t\int_0^\infty\partial_tP_{t-s}^{H}(x,y)f(s,y)dyds,\quad t,x\in(0,\infty),$$

for every $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$.

As in Lemma 2.1, we reduce the boundedness of $K\left(\{P_u^H\}_{u>0}\right)$ in $L^2((0,\infty)\times\mathbb{R},X)$ to study the operator $K_+\left(\{P_u^H\}_{u>0}\right)$ in $L^2((0,\infty)^2,X)$.

Lemma 4.1. *Let X be a Banach space. Then, the operator $K\left(\{P_u^H\}_{u>0}\right)$ is bounded in $L^2((0,\infty)\times\mathbb{R},X)$ if, and only if, $K_+\left(\{P_u^H\}_{u>0}\right)$ is bounded in $L^2((0,\infty)^2,X)$.*

Proof. Assume firstly that $K\left(\{P_u^H\}_{u>0}\right)$ is a bounded operator from $L^2((0,\infty)\times\mathbb{R},X)$ into itself, and let $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$. We define the extension f_0 of f as in (3). Then,

$$K_+\left(\{P_u^H\}_{u>0}\right)(f)(t,x)=K\left(\{P_u^H\}_{u>0}\right)(f_0)(t,x),\quad t,x\in(0,\infty),$$

and

$$\begin{aligned}\|K_+\left(\{P_u^H\}_{u>0}\right)(f)\|_{L^2((0,\infty)^2,X)}\leq&\|K\left(\{P_u^H\}_{u>0}\right)(f_0)\|_{L^2((0,\infty)\times\mathbb{R},X)}\\ \leq&\|f\|_{L^2((0,\infty)^2,X)}.\end{aligned}$$

Suppose now that $K_+\left(\{P_u^H\}_{u>0}\right)$ is bounded from $L^2((0,\infty)^2,X)$ into itself. Let $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$. We define operator $K_-\left(\{P_u^H\}_{u>0}\right)$ through

$$K_-\left(\{P_u^H\}_{u>0}\right)(f)(t,x)=\int_0^t\int_0^\infty\partial_tP_{t-s}^H(x,-y)f(s,y)dyds,\quad t,x\in(0,\infty).$$

We can write

$$\begin{aligned}\int_0^\infty\int_{\mathbb{R}}\left\|K\left(\{P_u^H\}_{u>0}\right)(f)(t,x)\right\|_X^2dxdt\leq 2\Big(&\int_0^\infty\int_0^\infty\left\|K_+\left(\{P_u^H\}_{u>0}\right)(f_+)(t,x)\right\|_X^2dxdt\\ &+\int_0^\infty\int_0^\infty\left\|K_-\left(\{P_u^H\}_{u>0}\right)(f_-)(t,x)\right\|_X^2dxdt+\int_0^\infty\int_0^\infty\left\|K_+\left(\{P_u^H\}_{u>0}\right)(f_-)(t,x)\right\|_X^2dxdt\\ &+\int_0^\infty\int_0^\infty\left\|K_-\left(\{P_u^H\}_{u>0}\right)(f_+)(t,x)\right\|_X^2dxdt\Big),\end{aligned}$$

where f_+ and f_- were defined in the proof of Lemma 2.1. In order to see that the operator $K\left(\{P_u^H\}_{u>0}\right)$ is bounded from $L^2((0,\infty)\times\mathbb{R},X)$ into itself, it is enough to prove that $K_-\left(\{P_u^H\}_{u>0}\right)$ is bounded in $L^2((0,\infty)^2,X)$.

We have that

$$\left|\partial_tP_{t-s}^H(x,-y)\right|\leq C\int_0^\infty\frac{e^{-c[|t-s|^2+(x+y)^2]/u}}{u^2}du\leq\frac{C}{|t-s|^2+(x+y)^2},\quad s,t,x,y\in(0,\infty).$$

Then, by proceeding as in the proof of Lemma 2.1 we conclude that $K_-\left(\{P_u^H\}_{u>0}\right)$ is bounded from $L^2((0,\infty)^2,X)$ into itself. □

We now split the operator $K_+\left(\{P_u^H\}_{u>0}\right)$ as follows

$$K_+\left(\{P_u^H\}_{u>0}\right)=K_+^{loc}\left(\{P_u^H\}_{u>0}\right)+K_+^{glob}\left(\{P_u^H\}_{u>0}\right),$$

where

$$K_+^{loc}\left(\{P_u^H\}_{u>0}\right)(f)(t,x)=\int_0^t\int_{x/2}^{2x}\partial_tP_{t-s}^H(x,y)f(s,y)dyds,\quad t,x\in(0,\infty),$$

for every $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$.

The operator $K(\{P_u^{L_\alpha}\}_{u>0})$ is also split in the following way

$$K\left(\{P_u^{L_\alpha}\}_{u>0}\right)=K^{loc}\left(\{P_u^{L_\alpha}\}_{u>0}\right)+K^{glob}\left(\{P_u^{L_\alpha}\}_{u>0}\right),$$

where

$$K^{loc}\left(\{P_u^{L_\alpha}\}_{u>0}\right)(f)(t,x)=\int_0^t\int_{x/2}^{2x}\partial_tP_{t-s}^{L_\alpha}(x,y)f(s,y)dyds,\quad t,x\in(0,\infty),$$

for every $f \in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$.

In the sequel we will use the following properties of the modified Bessel function I_ν, that can be encountered in [28, pp. 136 and 123]. The behavior of $I_\nu(z)$ close to zero is

$$I_\nu(z) \sim \frac{1}{2^\nu\Gamma(\nu+1)}z^\nu, \quad \text{as } z\to 0^+. \tag{11}$$

Moreover, for every $n\in\mathbb{N}$,

$$\sqrt{z}I_\nu(z) = \frac{e^z}{\sqrt{2\pi}}\left[\sum_{k=0}^n \frac{(-1)^k[\nu,k]}{(2z)^k} + \mathcal{O}\left(\frac{1}{z^{n+1}}\right)\right], \tag{12}$$

where $[\nu,0]=1$ and

$$[\nu,k] = \frac{(4\nu^2-1)(4\nu^2-3^2)\cdots(4\nu^2-(2k-1)^2)}{2^{2k}\Gamma(k+1)}, \quad k=1,2,\dots.$$

Lemma 4.2. *Let X be a Banach space. Then, the operators $K_+^{glob}(\{P_u^H\}_{u>0})$ and $K^{glob}\left(\{P_u^{L_\alpha}\}_{u>0}\right)$ are bounded in $L^2((0,\infty)\times\mathbb{R},X)$.*

Proof. Let $f\in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$. We have that

$$\left|\partial_t P_{t-s}^H(x,y)\right| \le C\int_0^\infty \frac{e^{-c[|t-s|^2+|x-y|^2]/u}}{u^2}du \le \frac{C}{|t-s|^2+|x-y|^2}, \quad s,t,x,y\in(0,\infty).$$

Hence, we obtain

$$\begin{aligned}\|K_+^{glob}(\{P_u^H\}_{u>0})(f)(t,x)\|_X &\le C\Big(\int_0^t\int_0^{x/2}\frac{1}{|t-s|^2+x^2}\|f(s,y)\|_X dyds \\ &\quad + \int_0^t\int_{2x}^\infty \frac{1}{|t-s|^2+y^2}\|f(s,y)\|_X dyds\Big) \\ &\le C\left(\frac{1}{x}\int_0^x P_*\left(\|\tilde f_0(\cdot,y)\|_X\right)(t)dy + \int_y^\infty \frac{1}{y}P_*\left(\|\tilde f_0(\cdot,y)\|_X\right)(t)dy\right), \quad t,x\in(0,\infty).\end{aligned}$$

Since the Hardy operators H_0 and H_∞ (see Section 2) are bounded in $L^2(0,\infty)$ and the maximal operator P_* is bounded in $L^2(\mathbb{R})$, we deduce that $K_+^{glob}(\{P_u^H\}_{u>0})$ is bounded in $L^2((0,\infty)^2,X)$.

According to (11) and (12) we get

$$0\le W_t^{L_\alpha}(x,y)\le C\frac{e^{-c|x-y|^2/t}}{\sqrt{t}}, \quad t,x,y\in(0,\infty).$$

By proceeding as above we obtain, for every $t,x\in(0,\infty)$,

$$\|K^{glob}(\{P_u^{L_\alpha}\}_{u>0})(f)(t,x)\|_X \le C\left(H_0\left[P_*\left(\|\tilde f_0(\cdot,y)\|_X\right)(t)\right](x) + H_\infty\left[P_*\left(\|\tilde f_0(\cdot,y)\|_X\right)(t)\right](x)\right).$$

Hence, $K^{glob}(\{P_u^{L_\alpha}\}_{u>0})$ is a bounded operator from $L^2((0,\infty)^2,X)$ into itself. □

From Lemmas 4.1 and 4.2 we deduce that the equivalence $(c)\Leftrightarrow(d)$ in Theorem 1.1 will be established once we prove the following.

Lemma 4.3. *Let X be a Banach space. Then, the operator $\mathbb{D}^{loc}$ given by*

$$\mathbb{D}^{loc} = K_+^{loc}\left(\{P_u^H\}_{u>0}\right) - K^{loc}\left(\{P_u^{L_\alpha}\}_{u>0}\right),$$

is bounded in $L^2((0,\infty)^2,X)$.

Proof. Let $f \in C_c^\infty(0,\infty)\otimes C_c^\infty(0,\infty)\otimes X$. We can write

$$\begin{aligned}\mathbb{D}^{loc}(f)(t,x) =&\frac{1}{\sqrt{4\pi}}\int_0^t\int_0^\infty \frac{e^{-|t-s|^2/4u}}{u^{3/2}}\left(1-\frac{|t-s|^2}{2u}\right)\\ &\times\int_{x/2}^{2x}\left[W_u^H(x,y)-W_u^{L_\alpha}(x,y)\right]f(s,y)dyduds, \quad t,x\in(0,\infty).\end{aligned}$$

By defining the function $\tilde{f}$ as in (6), we get for every $t,x\in(0,\infty)$,

$$\|\mathbb{D}^{loc}(f)(t,x)\|_X \le C\int_{x/2}^{2x} W_*(\|\tilde{f}(\cdot,y)\|_X)(t)\left(\int_0^\infty \left|W_u^H(x,y)-W_u^{L_\alpha}(x,y)\right|\frac{du}{u}\right)dy. \tag{13}$$

In order to estimate the function

$$H(x,y)=\int_0^\infty \left|W_u^H(x,y)-W_u^{L_\alpha}(x,y)\right|\frac{du}{u}, \quad x,y\in(0,\infty),\ x\neq y,$$

we distingue two cases. To simplify the notation we call $\xi=\xi(u,x,y)=2xye^{-u}/(1-e^{-2u})$, $u,x,y\in(0,\infty)$.

Suppose firstly that $\xi\ge 1$. By (8) and (12) we deduce

$$\begin{aligned}\left|W_u^H(x,y)-W_u^{L_\alpha}(x,y)\right| =&W_u^H(x,y)\left|1-\sqrt{2\pi}e^{-\xi}\sqrt{\xi}I_{\alpha-1/2}(\xi)\right|\le\frac{C}{\xi}W_u^H(x,y)\\ \le& C\begin{cases}\dfrac{e^{-c|x-y|^2/u}}{\xi^{1/4}\sqrt{u}}, & u\in(0,1),\\[2ex] \dfrac{e^{-u/2}e^{-c|x-y|^2}}{\xi^{1/4}}, & u\ge 1.\end{cases}\end{aligned}$$

Then,

$$\begin{aligned}\int_{0,\ \xi\ge1}^1 \left|W_u^H(x,y)-W_u^{L_\alpha}(x,y)\right|\frac{du}{u} \le& C\int_0^1\frac{e^{-c|x-y|^2/u}}{\xi^{1/4}}\frac{du}{u^{3/2}}\le\frac{C}{\sqrt{y}}\int_0^1\frac{e^{-c|x-y|^2/u}}{u^{5/4}}du\\ \le&\frac{C}{\sqrt{y|x-y|}}\int_0^\infty\frac{e^{-v}}{v^{3/4}}dv\le\frac{C}{y}\sqrt{\frac{y}{|x-y|}}, \quad \frac{x}{2}<y<2x,\ x\neq y.\end{aligned}\tag{14}$$

Also, we obtain

$$\begin{aligned}\int_{1,\ \xi\ge1}^\infty \left|W_u^H(x,y)-W_u^{L_\alpha}(x,y)\right|\frac{du}{u} \le& C\int_1^\infty\frac{e^{-u/2}e^{-c|x-y|^2}}{\xi^{1/4}}\frac{du}{u}\le\frac{e^{-c|x-y|^2}}{\sqrt{y}}\int_1^\infty e^{-u/4}du\\ \le&\frac{C}{y}\sqrt{\frac{y}{|x-y|}}, \quad \frac{x}{2}<y<2x,\ x\neq y.\end{aligned}\tag{15}$$

On the other hand, if $\xi\le 1$, by (11) we have that

$$\begin{aligned}\left|W_u^H(x,y)-W_u^{L_\alpha}(x,y)\right| \le& CW_u^H(x,y)\left(1+e^{-\xi}\sqrt{\xi}I_{\alpha-1/2}(\xi)\right)\le CW_u^H(x,y)\left(1+\xi^\alpha\right)\\ \le& C\begin{cases}\left(\dfrac{e^{-u}}{1-e^{-2u}}\right)^{1/4}\dfrac{e^{-c|x-y|^2/u}}{u^{1/4}}, & u\in(0,1),\\[2ex] \left(\dfrac{e^{-u}}{1-e^{-2u}}\right)^{1/4}e^{-cu}e^{-c|x-y|^2}, & u\ge 1.\end{cases}\end{aligned}$$

Then,

$$\begin{aligned}\int_{0,\ \xi\le1}^1 \left|W_u^H(x,y)-W_u^{L_\alpha}(x,y)\right|\frac{du}{u} \le& C\int_{0,\ \xi\le1}^1\frac{e^{-c|x-y|^2/u}}{u^{5/4}}\frac{\xi^{1/4}}{\sqrt{y}}du\\ \le&\frac{C}{\sqrt{y}}\int_0^1\frac{e^{-c|x-y|^2/u}}{u^{5/4}}du\le\frac{C}{y}\sqrt{\frac{y}{|x-y|}}, \quad \frac{x}{2}<y<2x,\ x\neq y,\end{aligned}\tag{16}$$

and

$$\begin{aligned}\int_{1,\ \xi\le 1}^{\infty}\left|W_u^H(x,y)-W_u^{L_\alpha}(x,y)\right|\frac{du}{u}\le& C\int_{1,\ \xi\le 1}^{\infty}\frac{\xi^{1/4}}{\sqrt{y}}e^{-cu}e^{-c|x-y|^2}du\\ \le&\frac{Ce^{-c|x-y|^2}}{\sqrt{y}}\int_1^\infty e^{-u/4}du\le\frac{C}{y}\sqrt{\frac{y}{|x-y|}},\quad \frac{x}{2}<y<2x,\ x\ne y. \qquad (17)\end{aligned}$$

From (13), (14), (15), (16) and (17) we deduce that

$$\|\mathbb{D}^{loc}(f)(t,x)\|_X\le C\mathcal{N}\left[W_*\left(\|\tilde{f}\|_X\right)(t)\right](x),\quad t,x\in(0,\infty), \qquad (18)$$

being

$$\mathcal{N}(f)=\int_{x/2}^{2x}\frac{1}{y}\sqrt{\frac{y}{|x-y|}}f(y)dy,\quad f\in L^2(0,\infty).$$

Jensen's inequality implies that the operator $\mathcal{N}$ is bounded from $L^2(0,\infty)$ into itself. Also, the maximal operator W_* is bounded in $L^2(\mathbb{R})$. Then, from (18) we infer that $\mathbb{D}^{loc}$ is a bounded operator from $L^2((0,\infty)^2,X)$ into itself. □

REFERENCES

[1] I. Abu-Falahah, P. R. Stinga, and J. L. Torrea, *Square functions associated to Schrödinger operators*, Studia Math., 203 (2011), pp. 171–194.

[2] I. Abu-Falahah and J. L. Torrea, *Hermite function expansions versus Hermite polynomial expansions*, Glasg. Math. J., 48 (2006), pp. 203–215.

[3] H. Amann, *Maximal regularity for nonautonomous evolution equations*, Adv. Nonlinear Stud., 4 (2004), pp. 417–430.

[4] W. Arendt and S. Bu, *The operator-valued Marcinkiewicz multiplier theorem and maximal regularity*, Math. Z., 240 (2002), pp. 311–343.

[5] ———, *Tools for maximal regularity*, Math. Proc. Cambridge Philos. Soc., 134 (2003), pp. 317–336.

[6] ———, *Fourier series in Banach spaces and maximal regularity*, in Vector measures, integration and related topics, vol. 201 of Oper. Theory Adv. Appl., Birkhäuser Verlag, Basel, 2010, pp. 21–39.

[7] P. Auscher and A. Axelsson, *Weighted maximal regularity estimates and solvability of non-smooth elliptic systems I*, Invent. Math., 184 (2011), pp. 47–115.

[8] P. Auscher, S. Monniaux, and P. Portal, *The maximal regularity operator on tent spaces*, Commun. Pure Appl. Anal., 11 (2012), pp. 2213–2219.

[9] J. J. Betancor, A. J. Castro, J. Curbelo, and L. Rodríguez-Mesa, *Characterization of UMD Banach spaces by imaginary powers of Hermite and Laguerre operators.* To appear in Complex Anal. Oper. Theory (`DOI: 10.1007/s11785-011-0203-9`).

[10] J. J. Betancor, A. J. Castro, and L. Rodríguez-Mesa, *Characterization of Banach valued BMO functions and UMD Banach spaces by using Bessel convolutions.* To appear in Positivity (`DOI: 10.1007/s11117-012-0189-1`).

[11] J. J. Betancor, A. Chicco Ruiz, J. C. Fariña, and L. Rodríguez-Mesa, *Odd* BMO($\mathbb{R}$) *functions and Carleson measures in the Bessel setting*, Integral Equations Operator Theory, 66 (2010), pp. 463–494.

[12] J. J. Betancor, R. Crescimbeni, J. C. Fariña, P. R. Stinga, and J. L. Torrea, *A* $T1$ *criterion for Hermite-Calderón-Zygmund operators on the* $BMO_H(\mathbb{R}^n)$ *space and applications.* To appear in Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) (`arXiv:1006.0416v2`).

[13] J. J. Betancor, J. C. Fariña, T. Martínez, and J. L. Torrea, *Riesz transform and g-function associated with Bessel operators and their appropriate Banach spaces*, Israel J. Math., 157 (2007), pp. 259–282.

[14] J. J. Betancor, J. C. Fariña, L. Rodríguez-Mesa, A. Sanabria, and J. L. Torrea, *Transference between Laguerre and Hermite settings*, J. Funct. Anal., 254 (2008), pp. 826–850.

[15] ———, *Lusin type and cotype for Laguerre g-functions*, Israel J. Math., 182 (2011), pp. 1–30.

[16] J. Bourgain, *Some remarks on Banach spaces in which martingale difference sequences are unconditional*, Ark. Mat., 21 (1983), pp. 163–168.

[17] D. L. Burkholder, *Martingales and Fourier analysis in Banach spaces*, in Probability and Analysis (Varenna, 1985), vol. 1206 of Lecture Notes in Math., Springer, Berlin, 1986, pp. 61–108.

[18] T. Coulhon and D. Lamberton, *Régularité L^p pour les équations d'évolution*, in Séminaire d'Analyse Fonctionelle 1984/1985, vol. 26 of Publ. Math. Univ. Paris VII, Paris, 1986, pp. 155–165.

[19] L. de Simon, *Un'applicazione della teoria degli integrali singolari allo studio delle equazioni differenziali lineari astratte del primo ordine*, Rend. Sem. Mat. Univ. Padova, 34 (1964), pp. 205–223.

[20] G. Dore and A. Venni, *On the closedness of the sum of two closed operators*, Math. Z., 196 (1987), pp. 189–201.

[21] J. Dziubański, G. Garrigós, T. Martínez, J. L. Torrea, and J. Zienkiewicz, *BMO spaces related to Schrödinger operators with potentials satisfying a reverse Hölder inequality*, Math. Z., 249 (2005), pp. 329–356.

[22] J. Dziubański and J. Zienkiewicz, *Hardy space H^1 associated to Schrödinger operator with potential satisfying reverse Hölder inequality*, Rev. Mat. Iberoamericana, 15 (1999), pp. 279–296.

[23] P. Grisvard, *Équations différentielles abstraites*, Ann. Sci. École Norm. Sup. (4), 2 (1969), pp. 311–395.

[24] S. Guerre-Delabrière, *Some remarks on complex powers of $(-\Delta)$ and UMD spaces*, Illinois J. Math., 35 (1991), pp. 401–407.

[25] E. Harboure, J. L. Torrea, and B. Viviani, *Vector-valued extensions of operators related to the Ornstein-Uhlenbeck semigroup*, J. Anal. Math., 91 (2003), pp. 1–29.

[26] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge, at the University Press, 1934.

[27] C. Le Merdy, *Counterexamples on L_p-maximal regularity*, Math. Z., 230 (1999), pp. 47–62.

[28] N. N. Lebedev, *Special functions and their applications*, Dover Publications Inc., New York, 1972.

[29] J. LeCrone, *Elliptic operators and maximal regularity on periodic little-Hölder spaces*, J. Evol. Equ., 12 (2012), pp. 295–325.

[30] B. Muckenhoupt, *Hermite conjugate expansions*, Trans. Amer. Math. Soc., 139 (1969), pp. 243–260.

[31] ———, *Poisson integrals for Hermite and Laguerre expansions*, Trans. Amer. Math. Soc., 139 (1969), pp. 231–242.

[32] ———, *Conjugate functions for Laguerre expansions*, Trans. Amer. Math. Soc., 147 (1970), pp. 403–418.

[33] B. Muckenhoupt and E. M. Stein, *Classical expansions and their relation to conjugate harmonic functions*, Trans. Amer. Math. Soc., 118 (1965), pp. 17–92.

[34] A. Nowak and K. Stempak, *Riesz transforms and conjugacy for Laguerre function expansions of Hermite type*, J. Funct. Anal., 244 (2007), pp. 399–443.

[35] ———, *Riesz transforms for multi-dimensional Laguerre function expansions*, Adv. Math., 215 (2007), pp. 642–678.

[36] J. L. Rubio de Francia, *Martingale and integral transforms of Banach space valued functions*, in Probability and Banach spaces (Zaragoza, 1985), vol. 1221 of Lecture Notes in Math., Springer, Berlin, 1986, pp. 195–222.

[37] K. Stempak and J. L. Torrea, *Poisson integrals and Riesz transforms for Hermite function expansions with weights*, J. Funct. Anal., 202 (2003), pp. 443–472.

[38] ———, *On g-functions for Hermite function expansions*, Acta Math. Hungar., 109 (2005), pp. 99–125.

[39] G. Szegő, *Orthogonal polynomials*, American Mathematical Society, Providence, R.I., fourth ed., 1975. American Mathematical Society, Colloquium Publications, Vol. XXIII.

[40] S. Thangavelu, *Lectures on Hermite and Laguerre expansions*, vol. 42 of Mathematical Notes, Princeton University Press, Princeton, NJ, 1993.

Víctor Almeida, Jorge J. Betancor, Alejandro J. Castro
Departamento de Análisis Matemático, Universidad de La Laguna,
Campus de Anchieta, Avda. Astrofísico Francisco Sánchez, s/n,
38271, La Laguna (Sta. Cruz de Tenerife), Spain
E-mail address: `valmeida@ull.es, jbetanco@ull.es, ajcastro@ull.es`